



## Which foodweb?


- Freshwater Delta
- Low-Salinity Zone (LSZ, ~ 0.5 5 or 10)
- Salty regions South Bay-San Pablo Bay

## Main points


- 1. Plankton are quasi-Lagrangian particles
- 2. Limits to primary production
- 3. Consequent limits to zooplankton production
- 4. Spatial subsidies
- 5. How to design an estuarine copepod
- 6. Can we fix the low productivity?
- 7. Some key questions

Some old stuff, some new!


#### 1. Plankton are quasi-Lagrangian particles



#### 1. Plankton are quasi-Lagrangian particles



#### Effect of particle behavior on vertical position



**Depth as Fraction of Water Column** 


1.0

1128 particles (max)

Along-Channel Velocity, m s<sup>-1</sup>

Kimmerer Gross MacWilliams In Press Limnol. Oceanogr.

## Effect of particle behavior on vertical position




Areas of circles ∞ number of particles

1128 particles (max)

Kimmerer Gross MacWIlliams In Press Limnol. Oceanogr.

## Effect of particle behavior on transport

Tidal Biased ↓ on ebb ↑ on flood



Proportion of Particles

Lost at Sea

>10

5 - 10


0.5 - 5.0

0.0 - 0.5

Days since Release

## Effect of particle behavior on transport





Proportion of Particles

Lost at Sea

>10


5 - 10

0.5 - 5.0

0.0 - 0.5


Days since Release

#### 2. Limits to primary production: light




Primary Productivity (mgC m<sup>-3</sup>)

#### 2. Limits to primary production: Grazing




### 2. Limits to primary production: Grazing




Kimmerer & Thompson 2014 Estuaries & Coasts

#### 2. Limits to primary production: Size fractions

111 paired sets of <sup>14</sup>C incubations 5 µm and GF/F filter




#### 2. Limits to primary production: Net growth 1987-2008




2. Limits to primary production: Ammonium inhibition

Ammonium Inhibition

## 2. Limits to primary production: Conceptual Model



#### 2. Limits to primary production: Conceptual Model



<sup>14</sup>C Measurements

Mixing diagrams

Growth rate & efficiency


**Dilution experiments** 

Nutrient bioassays


Clam biomass and filtration rate (spatially intensive surveys)

Symbols: Integration and Application Network, UMD Center for Environmental Science


#### 3. Limits to zooplankton production: Low-salinity foodweb



#### 3. Limits to zooplankton production: Foodweb



#### 3. Limits to zooplankton production: slow development




Kimmerer et al. Fall Habitat study

#### 3. Limits to zooplankton production: slow development

# Development Rate ∞ Phytoplankton Biomass







Sac SJR

- Freshwater
- Low Salinity


### 3. Limits to zooplankton production: predation






York et al. 2013 Estuaries & Coasts & Slaughter et al. in prep.


## 4. Spatial subsidies: Conceptual model




## 4. Spatial subsidies: Conceptual model



## 4. Spatial subsidies: Conceptual model




## 4. Spatial subsidies: phytoplankton



2014 Estuaries & Coasts


## 4. Spatial subsidies: phytoplankton



## 4. Spatial subsidies: copepod distributions




Pseudodiaptomus forbesi



Fall Habitat study

## 4. Spatial subsidies: copepods in the Low-Salinity Zone






Pseudodiaptomus forbesi

## 5. How to design an estuarine copepod: Method 1

- Phytoplankton are small
- Low food supply
- Most fish are visual predators
- Few filter-feeding predators

Eat ciliates
Low feeding rate
Small, cryptic
Slow growth OK





## 5. How to design an estuarine copepod: Method 2


- Phytoplankton are small
- Low food supply
- Most fish are visual predators
- Few filter-feeding predators

Don't eat 'em

Eat copepods


**Grow fast** 

Doesn't matter





Acartiella sinensis



Slaughter et al. Fall Habitat study

#### 5. How to design an estuarine copepod: Method 3

- Phytoplankton are small
- Low food supply
- Most fish are visual predators
- Few filter-feeding predators

Be omnivorous


??

Spatial refuge

Doesn't matter



Pseudodiaptomus forbesi



Adults

Copepodites


Nauplii

Kimmerer et al. Fall Habitat study

### 6. Can we fix the low productivity?


- Alter productivity directly
- Increase the large-scale spatial subsidy
- Use small-scale spatial subsidies
  - Marsh restoration

#### 6. Can we fix the low productivity?



Ammonium? Clams? Turbidity?

#### 6. Can we fix the low productivity? Large-scale export




High flow increases subsidy to Low-Salinity zone



But it costs a bundle!

3 months @ \$0.10 / m<sup>3</sup> or \$120/AF

#### 6. Can we fix the low productivity? Large-scale export




High flow increases subsidy to Low-Salinity zone



Total capacity of major Sacramento Valley reservoirs

### 6. Can we fix the low productivity? Marsh export



Shallow water:
High growth
Benthic grazing
Habitat for planktivores


Production:
Total organic matter
Phytoplankton
Zooplankton


++

±

-'?

#### 6. Can we fix the low productivity? Marsh export






#### Open Water

How much stimulation
by marsh production?
Phytoplankton
Zooplankton
Detritus
Nekton

### 6. Can we fix the low productivity? Export of plankton



#### **Marsh**

Area 1000 ha

Depth 2m

Copepods 23 mgC m<sup>-3</sup>

Growth rate  $\mu$  0.1 d<sup>-1</sup>

Residence time 10d

#### **Open Water**

Volume 0.5 km<sup>3</sup>

Copepods 3 mgC m<sup>-3</sup>

Resulting subsidy: 3% of existing copepod biomass d<sup>-1</sup>




#### 6. Can we fix the low productivity? Literature review


## Flux of zooplankton to and from marshes

- Lake zooplankton abundance low in shoals
- Marsh in NY a sink for zooplankton
- Australian tidal creek a sink for copepods
- Saltmarsh a sink for copepods, source for larvae
- China Camp: marsh a sink for mysids
- Liberty Island: source or sink


### 7. Some key questions: marsh subsidy




### 7. Some key questions: marsh subsidy




#### 7. Some key questions: phytoplankton production



#### 7. Some key questions: phytoplankton production



#### 7. Some key questions: phytoplankton production




# Summary

- Plankton move with the water
  - Use their frame of reference
- Three reasons for low foodweb productivity
  - Light limitation
  - Grazing
  - Ammonium concentration
- Three unlikely remedies
  - Increase outflow
  - Get rid of clams
  - Plankton farms
- Three potential remedies
  - Reduce ammonium loading
  - Marshes as plankton factories
  - Detrital output of marshes

Will this be science-based, or faith-based?

# Summary

- Plankton move with the water
  - Use their frame of reference
- Three reasons for low foodweb productivity
  - Light limitation
  - Grazing
  - Ammonium concentration



#### Thanks to:


Anne Slaughter, Toni Ignoffo, Lindsay Sullivan, Karen Kayfetz, Jan Thompson, April Hennessy, Ed Gross, Michael MacWilliams, Erwin Van Nieuwenhuyse,

. . .



Funding from
Delta Science Program
Interagency Ecological Program
National Science Foundation

## Subsidies from marsh: phyto model



Area
Depth
Phytoplankton
Growth rate 

Microzoo grazing
Residence time


1000 ha 2m 900 mgC m<sup>-3</sup> 0.86 d<sup>-1</sup> 60% μ 10d Volume Phytoplankton

0.5 km<sup>3</sup> 73 mgC m<sup>-3</sup>



Resulting subsidy: 5% of existing phytoplankton biomass d<sup>-1</sup>

#### Lesson 9: Foodweb interactions can be subtle



# Zooplankton as "ecological filters" \*

How does variation in phytoplankton productivity translate to variability in fish recruitment?

	Efficiency of transfer to fish	
Characteristic	Low	High
Phytoplankton size	Small	Large
Seasonal pattern (vs. fish)	Unpredictable	Predictable
Sensitivity to food	Low	High
Reproductive mode	Egg carriers	Broadcast spawners
Palatability / nutrition	Low	High
Zooplankton size (fish size)	Small	Large
Detectability (feeding mode)	Low	High
Catchability (feeding mode)	Low	High
Foodweb length	Long	Short